

EFFECT OF IONISED (ELECTROLYSED) WATER ON THE RAT EMBRYO DEVELOPMENT

Jolita STANKEVIČ¹, Aistė AUDICKAITĖ¹, Simona ŠILOVĖ^{1*}, Valdas ŠIMČIKAS²,
Henrikas CESIULIS³, Grita SKUJIENĖ⁴, Virginija BUKELSKIENĖ⁵,
Violeta ŽALGEVIČIENĖ¹, Janina TUTKUVIENĖ¹

¹Department of Anatomy, Histology and Anthropology of the Institute of Biomedical Sciences,
Faculty of Medicine, Vilnius University, Čiurlionio g. 21/27, LT-03101, Vilnius, Lithuania

²Joint Stock Company “Water research”, Žirmūnų g. 106, LT-09121, Vilnius, Lithuania

³Department of Physical Chemistry of the Faculty of Chemistry and Geosciences, Vilnius University,
Naugarduko g. 24, LT-03225, Vilnius, Lithuania

⁴Department of Zoology of the Institute of Biosciences, Life Sciences Center, Vilnius University,
Saulėtekio al. 7, LT-10223, Vilnius, Lithuania

⁵Department of Biological Models of the Institute of Biochemistry, Life Sciences Center,
Vilnius University, Saulėtekio al. 7, LT-10223, Vilnius, Lithuania

Received 05 June 2019; accepted 18 November 2019

Abstract. The aim of this study was to investigate the effects ionised water has on embryonic development using Wistar rat animal model. For that purpose, alkaline and acidic water was prepared with a domestic water ioniser. It was found that the concentrations of Cl^- , SO_4^{2-} ions increased in acidic water, while in alkaline water, Ca^{2+} concentration decreased and halogenated hydrocarbon concentrations exceeded permitted levels. The animals were given test alkaline and acidic water, as well as tap water as control. After three months, female rats were mated. On the 21st day of gestation, they were euthanized and subjected to Caesarean sections; the number of live and dead fetuses was recorded. The fetuses were examined for external or visceral malformations and skeletal abnormalities. The data showed that embryo death was higher in acidic and alkaline experimental groups in comparison to the control group. The fetuses in both test groups were significantly shorter than in the control group. Long bones of fetal hind and front limbs were shorter in the acidic group in comparison to the control group. Retardation of limb osteogenesis was expressed in the acidic group fetuses. Therefore, in our model, ionised water had a negative effect on the embryonic development.

Keywords: alkaline and acidic water, electrolysis, rat, embryonic development, water cleaning technologies.

Introduction

Water is the most important substance, which determines survival of all living beings. In the organism, it has a number of roles: it serves as a solvent, temperature buffer, metabolite, living environment, etc. However, natural water is not chemically pure H_2O . Atmospheric precipitation, rocks and sediments, through which water is filtered, play a key role in the formation of underground freshwater. Therefore, the variety of chemical compounds, found in drinking water, depends on the chemical composition of the rock (Lack, 1999). Consumption of water also determines the supply of any minerals it contains. Research suggests that 14 out of 21 mineral elements required for

humans are essential for good health. Different combinations of these elements are responsible for managing normal functioning of the body, i.e. bone and membrane structure (Ca, P, Mg, F), water and electrolyte balance (Na, K, Cl), metabolic catalysis (Zn, Cu, Se, Mg, Mn, Mo), oxygen binding (Fe), and hormone functions (I, Cr) (World Health Organization, 2005). However, though most of the chemical elements, present in water, are required for a healthy functioning of a human body, surplus in these elements may lead to various diseases and functional problems; thus, the amount of these elements allowed in drinking water is regulated and limited.

Lithuania is considered among the leaders in EU in terms of water purity (LRP, 2016). Specialists claim that

*Corresponding author. E-mail: simona.silove@gmail.com

unpolluted deep water aquifers in Lithuania can be envied by most of the countries for groundwater quality, chemical composition and microbiological indicators. The water is not contaminated with such trace elements as iodine, while others, e.g. fluoride, are very unevenly distributed throughout the country (Klimas & Mališauskas, 2008). Nitrate, commonly associated with agricultural activities, is also found in very small quantities in Lithuanian groundwater. The same applies to other north European countries, such as Norway, Sweden, Finland, Estonia, and Latvia. It is believed that this is due to frozen ground in winter time, when the land is not arable, thus cannot be polluted with fertilisers and pesticides, the surplus of which might seep into the groundwater (Flem, Reimann, Birke, Banks, & Filzmoser, 2015).

Moreover, in Europe both surface (from rivers or lakes) and ground water is used for public water supply, while in some countries (Lithuania, Denmark, and Austria) mostly underground water is used for drinking (Volker & Borchardt, 2019; Flem et al., 2015). Most Lithuanian waters require no treatment, while those requiring some form of purification, are treated with sodium hypochlorite solution, which is considered safe for the environment and people (Vilniaus vandenys, 2016). Still, popular literature and the Internet abound with articles and advertising projects that recommend water treatment with water ionisers, which through the process of electrolysis produce two types of water: acidic (dead) and alkaline (living) (Laucevičius, 2009).

It is known that naturally, with the help of the solar energy, electrolysis occurs in plants and algae. In industry, electrolysis is performed to clean and preserve metallic objects from corrosion, while in the future even hydrogen might be produced by electrolysis as an alternative to fossil fuel (Santos et al., 2019; Rashid, Al Mesfer, Naseem, & Danish, 2015). Meanwhile, some people believe that electrolysis makes water better and that drinking it may cure many different illnesses. The somewhat popular alkaline diet adds to this believe. However, according to Fenton and Huang (2016), there is no proof that alkaline diet or alkaline water, as promoted by sales agents selling water alkalinisers, has any effect on cancer prevention or treatment.

Still, authors who have been experimenting with living organisms in the hope that alkaline ionised water can work as a protection against diseases stemming from oxidative stress – diabetes, cancer, arteriosclerosis, neurodegenerative diseases, and adverse effects of hemodialysis – found that alkaline water is actually beneficial for human health (Xue et al., 2014; Ignacio, Joo, & Lee, 2012; Shirahata, Hamasaki, & Teruya, 2012). In addition, some authors brought acid-base hypothesis into view, which suggests that an overly acidic diet (high in protein and grains) leads to chronic diseases and bone health problems, thus, raising the body pH through alkaline diet and balancing the acid-base ratio could prevent osteoporosis and other related diseases (Dawson-Hughes, 2016; Xu et al., 2016; Fenton, Eliasziw, Lyon, Tough, & Hanley, 2008). However, as the authors state, the results on acid-base hypothesis are

inconclusive and require further investigations (Dawson-Hughes, 2016; C. J. Fenton, T. R. Fenton, & Huang, 2017; Fenton et al., 2008)

In nature, high levels of either acidity or alkalinity can destroy life. Acid rains affect the soil, which in turn becomes less fertile and has a negative effect on microorganisms, flora and fauna living on it. Likewise, contaminants responsible for air pollution, such as nitrogen oxides and sulphur dioxides – waste product of manufacturing, transportation, mining and agriculture – when introduced into water, transform into acids and significantly affect water's pH levels (Osman, 2018; Weldelessie, Naz, Singh, & Oves, 2018). In the presence of such information, the authors of this study found it important to investigate the effect electrolysis-altered water has on living organisms.

It is known that during electrolysis, ions distribute unevenly in the solution. Anions Cl^- , SO_4^{2-} , NO_3^- migrate towards the anode, where oxidation reactions take place, producing free chlorine and oxygen gas. Depending on the composition of the electrolysed solution, various salts, sulphuric or nitric acids form, and the pH value falls below 7. Meanwhile, cations migrate towards the cathode, where reduction occurs. Depending on the alkalinity of the solution, Ca^{2+} and Mg^{2+} ions might form low solubility hydroxides (resulting in turbidity) whereas sodium and potassium stay in the solution and form soluble sodium and potassium hydroxides. Potassium hydroxide begins to dominate the solution, and the pH value increases to 8–9.

Normally, halogenated hydrocarbons are not found in nature, as they are a product of human activity. In water, halogenated hydrocarbons can form after chlorination or by other means of water disinfection or electrolysis. Free chlorine, which forms during electrolysis, reacts with water, bromide, and organic substances, resulting in a variety of halogenated compounds – usually haloforms. Concentrations of free chlorine and haloforms are directly dependent on the composition of water being electrolysed (especially the initial concentration of Cl^-), as well as on the duration of electrolysis.

According to the proponents of water ionisers, acidic water produced in these devices is a natural bactericide and pesticide (Bui et al., 2017; Sun, Zhang, Chen, & Han, 2012; Abbasi & Lazarovits, 2006). They also claim that alkaline water is better in neutralising stomach gastric acid, detoxifies the organism, and acts like a natural antioxidant and alkalizing substance due to a large content of oxygen and negative oxidation-reduction potential (Mousa, 2016; Laucevičius, 2009). There are many popular and scientific articles about the benefits of electrolysed water (Weidman et al., 2016; Koufman & Johnston, 2012; Shirahata et al., 2012; Laucevičius, 2009; Shirahata et al., 1997). Because of that, significant part of society now claims to have found a surprising “drug” for treating many different ailments. However, there is little research on the adverse effects electrolysed water has on living organisms (El-Fiky, 2002; Merne, K. J. Syrjanen, & S. M. Syrjanen, 2001; Watanabe & Kishikawa, 1998a; Watanabe et al., 1998b; Watanabe, Kishikawa, & Shirai, 1997).

Thus, it can be argued that there is a lack of aggregated studies on electrolysed water effect on the whole organism. Likewise, not much is known how electrolysed water affects the development of offspring. Therefore, due to deficiency of scientific evidence supporting the safety of electrolysed water, and due to its popularity in Lithuania and other countries, the aim of this study was to assess the effects of ionised (electrolysed) water on the embryonic development using Wistar rats as a test model. Our goal was to evaluate the formation of the offspring of female rats, which were given acidic and alkaline water before and during pregnancy.

1. Materials and methods

1.1. Water electrolysis

Tap water provided by "Vilniaus vandenys" Ltd (Lithuania) was used in this study. To produce alkaline and acidic water, ordinary tap water was treated with a domestic water ioniser ("PTV-KL") for 10 and 20 min respectively, according to manufacturer's instructions.

1.2. Measurement of main ions and volatile halogenated hydrocarbons in the water

The ion analysis was carried out using Ion Chromatograph DIONEX IC-1000 according to Standards ISO 14911 (Cations) and ISO 10304-1 (Anions).

Haloform analysis was carried out by Gas Chromatograph DANI GC 1000 using Head Space injection technique according to Standard ISO 10301. The samples were taken from sealed vials in which the ratio of water volume to air volume was fixed. In this study, 20 mL-sized vials were filled with 10 mL of samples. The temperature of the vials was stabilized in a thermostatic system at 70 °C to achieve specified equilibrium conditions. After reaching equilibrium with water, an electron capture detector was used for gas chromatography in the sampling vials.

1.3. Animals

In this study, 3 months old Wistar rats were used. Approval of Ethics Committee and permission for the experimentation was received from the State Food and Veterinary Service of Lithuania, No 1; 14-01-2013. The experiment was designed in accordance to the requirements stated in 2010/63/EU Directive and Order of the Lithuanian State Food and Veterinary Service Director No B1-866; 31-12-2012. Wistar rats were obtained from animal experiment was carried out in the Department of Biological Models, Vilnius University, Institute of Biochemistry (Lithuania).

1.4. Experimental methodology

Three months before mating, female rats were divided into three groups and given different types of electrolysed water: alkaline ($n = 10$) or acidic ($n = 10$), and tap water

as control ($n = 10$). During this period and at the time of gestation, rats were provided with standard food for rodents and electrolysed (or tap) water *ad libitum*. The animals were weighed regularly.

After three months, female Wistar rats were mated overnight with males of the same clone. The following morning, vaginal smears were subjected to a microscopic examination in order to determine the presence of sperm. The day of sperm detection in vaginal smears was designated as day 0 of gestation. On the 21st day of gestation, the animals were euthanized and subjected to Caesarean sections. The uteri were removed and opened; dead and live fetuses were recorded in the uterine horn to determine post-implantation mortality indices.

1.5. Measurement of mother rat blood open circuit potential

Values of open circuit potential (OCP) were calculated from recorded time dependencies of potential differences measured directly between platinum electrode and reference electrode (saturated Ag, AgCl/Cl⁻ electrode) immersed into the sample of blood. The volume of the sample was ~1 mL. The dependencies were recorded by means of programmable potentiostat/galvanostat AUTOLAB N302. Heparin was used as the anticoagulant of blood.

1.6. Analysis of fetuses

To assess the effects of ionised (electrolysed) water on the embryonic development, external and internal physical features of the fetuses were evaluated. For that purpose, the fetuses were weighed and measured; half fetuses of every female (alkaline ($n = 35$), acidic ($n = 20$), control ($n = 47$)) were fixed in Bouin's solution for subsequent examination for external and visceral malformations. In order to render the skeleton visible, another set of fetuses (alkaline ($n = 34$), acidic ($n = 21$), control ($n = 48$)) was fixed in ethanol. For visceral malformation analysis, fetuses were sliced using Hayes technique. Using an eyepiece micrometre attached to a stereomicroscope (MBS-1), internal organs of the fetuses (liver, kidney, heart, lungs, brain and other) were investigated, and length of ossification centers of the fetal limb bones was measured in the middle of diaphysis of long tubular bones (Hayes, 1994). The fetuses of the control group were prepared and examined in the same way as the experimental ones.

1.7. Statistical analysis

All the data are expressed as mean \pm standard deviation (SD). Two-tailed Student's t-test was performed to ascertain statistical significance ($p < 0.05$) in differences between results of experimental and control groups. For the evaluation of ion concentrations in the water and water's effect on bone length, one-way analysis of variance (one way-ANOVA) was used, followed by Tukey's multiple comparisons post-hoc.

2. Results

2.1. Ion concentration in water and water pH after electrolysis

In the experiments, after tap water electrolysis for 10 and 20 min, concentration of main ions, volatile halogenated hydrocarbons and pH were analysed. The results revealed, that pH of electrolysed water assay was significantly different from the tap water (pH – 7.91) used: pH of acidic water was 3.38, and of alkaline – 10.14. Also, the concentration of Ca²⁺ was significantly reduced in alkaline water as compared to tap water (Figure 1a and 1b).

The study of halogenated hydrocarbons in electrolysed water demonstrated increased concentrations of chloroform (CHCl₃) and bromodichloromethane (CHBrCl₂) in both acidic and alkaline water (Table 1). After ionisation, the concentration of chloroform in alkaline water increased from 0.38 (after 10 min) to 0.63 µg/l (after 20 min), while in acidic water, it decreased from 25.20 (after 10 min) to 23.10 µg/l (after 20 min). Similarly, the concentration of bromodichloromethane increased in acidic water from 10.50 to 16.30 µg/l. Also, an increase of bromoforms was detected in acidic water – from 0.89 to 1.49 µg/l.

2.2. Blood OCP, blood and urine pH in mother rats

In the medical, biological, and ecological literature, an OCP of platinum electrode placed into a test medium against a certain reference electrode is considered as the redox potential of the medium. In this case, “redox potential” means the OCP of the electrode immersed into studied biological medium, such as blood or urine.

To evaluate the impact of ionised water on the physical status of mother rats, blood and urine samples were collected just before mating. It was found that for rats in acidic and alkaline groups, the values of OCP in blood decreased by ~0.03 V: from 0.165 V in control group to 0.135 V in groups, which consumed ionised water.

Furthermore, urine pH of rats, which were given electrolysed water for 3 months, ranged as follows: 7.59±0.40 in acidic, and 7.23±0.53 in alkaline groups. Urine pH of control rats, which were given tap water, was 7.35±0.58. Therefore, generally urine pH did not differ significantly between the experimental and control groups (p > 0.05).




Figure 1. Ion (Cl⁻, SO₄²⁻, NO₃⁻, Na⁺, K⁺, Ca²⁺, Mg²⁺) concentrations in acidic, alkaline and control tap water (mg/l); a – after 10 min.; b – after 20 min. of electrolysis; mean ± S.D.; *p < 0.05 when compared to control; #p < 0.05 when compared to alkaline group

Similarly, mother rats’ blood pH did not differ in experimental and control groups, when compared (p > 0.05): blood pH of alkaline group was 7.78±0.17, acidic group – 7.80±0.17, and control group – 7.80±0.10. Thus, the data suggests that ionised water did not affect the acidity or basicity of experimental rats’ blood.

2.3. Body weight of differently watered rats

The analysis of rats’ body weight gained during three months revealed such results: animals given alkaline water gained 40.5±18.05 g, acidic water – 32.4±16.90 g, while during the same time rats in the control group gained 52.4±17.52 g. Thus, rats in both experimental groups gained less weight than those in the control group, the lowest weight gain being observed in the acidic group. Nonetheless, there were no statistically significant

Table 1. Dissolved halogenated hydrocarbons (chloroform, bromodichloromethane, chlorodibromomethane, bromoform) in control (tap water), alkaline and acidic water after 10 and 20 min of electrolysis (µg/l)

		Chloroform (µg/l)	Bromodichloromethane (µg/l)	Chlorodibromomethane (µg/l)	Bromoform (µg/l)
Control	Tap water	<0.1	<0.1	<0.1	<0.1
Alkaline water	10 min electrolysis	0.38±0.02	0.13±0.04	<0.1	<0.1
	20 min electrolysis	0.63±0.16	0.13±0.01	<0.1	<0.1
Acidic water	10 min electrolysis	25.20±0.99	10.50±2.84	5.04±1.38	0.89±0.11
	20 min electrolysis	23.10±6.08	16.30±4.66	12.40±1.41	1.49±0.01




Figure 2. Ionised water effect on fetus length:

a – general view of control (left) and acidic (right) groups fetuses; b – fetuses of control (left) and acidic (right) groups represent the retardation of bone development in acidic group fetuses; c – comparison of fetuses' length in control, alkaline and acidic groups; acidic group fetuses were significantly shorter ($p < 0.05$) compared to control

differences between both experimental groups and control group ($p > 0.05$).

It also should be noted that rats in both experimental groups tended to consume less water (acidic or alkaline) than rats in control group, which consumed tap water.

2.4. Embryonic development

The main objective of this study was to assess the effects of ionised (electrolysed) water taken by the mother rats on the embryonic development. In assessing the viability of rat embryos, the data of this study revealed that embryo death rate increased in both acidic (27.0%) and alkaline (10.1%) groups compared to the control group (0%). In addition, it was detected that female rats, which had been given ionised water, had 10.14% of embryo resorptions. On the contrary, no embryo resorption was detected in rats, which consumed tap water.

Moreover, electrolysed water did not cause any noticeable developmental defects of external or internal organs of the fetuses (Figure 2a and 2b). However, ionised water had a noticeable effect on the size of fetuses: rat fetuses of both experimental groups were shorter than those of the control group; even more, acidic group fetuses were significantly shorter ($p < 0.05$) compared to the controls (Figure 2c), even though their weight did not differ.

While evaluating ossification centers of front and hind limb bones in coloured skeletons of the fetuses, it was found that acidic water did influence ossification of the long bones of limbs – they were shorter in comparison to the control group ($p < 0.05$) (Figure 3a and 3b). On the

other hand, ossification of the long bones of the fetuses from alkaline group had no disruptions in skeletal development, and was not statistically different from the fetuses in the control group ($p > 0.05$).

3. Discussion

Water ionisation supporters claim that alkaline water neutralizes bodily acids and helps with the prevention of diseases, however, according to Vormann and Goedecke (2002), various buffer systems are responsible for a stable pH of body fluids, thus the acidity or alkalinity of food and drink is irrelevant, as it will not change the pH values of the body. In agreement with that, the results of this study indicate that urine and blood pH of the rats given alkaline water did not become more alkaline. Moreover, female rats which were given acidic and alkaline water gained less weight, though no statistically significant difference between experimental and control groups was found. These results could be compared to those of Watanabe et al. (Watanabe & Kishikawa, 1998a; Watanabe et al., 1998b; Watanabe et al., 1997), who investigated how alkaline water changed body weight, and analysed biochemical parameters of the blood, as well as hearts of pregnant rats and their offspring. Their study demonstrated that alkaline water leads to increased body weight of rats and some of their organs, erythrocyte haemolysis,




Figure 3. Effect of electrolysed water on the ossification of the fetuses long bones of limbs: a – length of front leg bud in control, alkaline and acidic group fetuses; b – length of hind leg bud in control, alkaline and acidic group fetuses; mean \pm S.D.; * $p < 0.05$ when compared to control; # $p < 0.05$ when compared to acidic group

elevated blood potassium, and induced myocardial necrosis and fibrosis.

On the other hand, Massimiliano Magro's et al. (2016) 3-year survival study of alkaline water consumption effects on a population of mice revealed that even though animals which were given alkaline water lived longer, there was no correlation between a risk of diseases and alkaline water consumption. After examining the animals' organs histologically, the authors found that there were no significant differences between mice treated with alkaline water and tap water (Magro et al., 2016). However, it must be taken into consideration that water ionisers produce both alkaline and acidic water, and that there is a permeable membrane between the cameras of the device, where this water is produced.

During electrolysis halogenated hydrocarbons (chloroform, bromodichloromethane, chlorodibromoethane, and bromoform) increase in concentration in the water. Out of these, the levels of chloroform are usually the most elevated. Through the aforementioned membrane, part of the chlorine ions migrate from acidic to alkaline water. It is well known that chloroform is used for water disinfection and that when its concentration increases in water, together with bromodichloromethane, it may become carcinogenic and increase the risk of malignancies – colon cancer, premature births or embryonic anomalies (Benmarhnia, Delpla, Schwarz, Rodriguez, & Levallois, 2018; Abdelhalim, Salaheldeen, Idris, Abdelsalam, & Sabahelkhier, 2016; Hrudey et al., 2015; Villanueva, Cordier, Font-Ribera, Salas, & Levallois, 2015; Chen, Lin, Duh, Chou, & Hsu, 2011; Florentin, Hautemaniere, & Hartemann, 2011; Legay et al., 2011; Mills et al., 1998). Our study revealed that even after 10 minutes of water electrolysis, halogenated compound levels of methane and ethane (haloforms) – toxic for living organisms – increased in the water. Thus, even though in the present study electrolysed water did not cause any noticeable developmental defects of the external or internal organs of the fetuses, nevertheless, this water had a significant impact on fetus size. Significant differences between ossification centers in long bones of fetuses were also detected.

Various literature sources point out that embryonic ossification depends on genetics or is affected by oxidative stress, diet, hormones, various environmental factors, and maternal physical activity (Hu et al., 2018; Jensen et al., 2017; Fadel, Sequeira, Abu-Hijleh, Obeidat, & Salem, 2012; Ornoy, Rand, & Bischitz, 2010; Rauch & Schoenau, 2001; Fadel & Persaud, 1993). Likewise, limb ossification disorders can be induced by teratogenic substances, such as azathioprine, which impairs the interaction between the apical ectodermal ridge and the progress zone, i.e., the morphogenetic zones located in the distal part of the limb (Zukiene, Zalgeviene, & Rizgeliene, 2003). The results of this study demonstrated that, prior to mating, mother rats, which were given acidic water, tended to weigh less. Stress indicators, such as rough coat and bloody tears, were also evident in this animal group. Such physical changes of

mother rats could have been the cause of embryo growth retardation and the slowing down of their bone development. In addition, reduced Ca^{2+} concentration in ionised water could also have led to abnormal ossification. Also, the possibility that halogenated hydrocarbons detected in water after electrolysis could be the cause of ossification retardation and other ossification disorders in the embryo must equally be taken into consideration.

Furthermore, in medical science, many attempts were made to estimate the redox properties of blood, other biological liquids, and tissues by measuring OCP. In a previous study done by the authors, rat blood OCP values decreased by ~ 0.03 V for rats in acidic and alkaline groups in comparison to the control group (Audickaitė et al., 2014). Usually, such values indicate no presence of pathology in human blood (Khubutiya et al., 2010). However, if pathology was detected, notable deviations from the average values of OCP would appear, and differences between OCP could reach up to 0.1 V (Vormann & Goedecke, 2002). Since many homeostatic processes are electrochemical, when their activity increases, and changes in the composition of potential forming system appear, OCP levels vary more significantly. However, evaluation according to the Nernst equation reveals that the ratio of oxidized molecules to reduced molecules in the rats' blood decreases up to 3 times, and such a decrease is probably enough to cause pathology in rats.

Conclusions

1. The study revealed that after electrolysis the concentrations of Cl^- , SO_4^{2-} ions increased in acidic water, while in alkaline water Ca^{2+} concentration decreased and halogenated hydrocarbon concentrations exceeded permitted levels.
2. Rats, which were given alkaline and acidic water for 3 months, tended to gain less weight in comparison to those that drank tap water.
3. Blood and urine pH differences were not statistically significant between groups; however, the urine of rats, which had been given alkaline water, had a tendency to acidify.
4. Neither alkaline nor acidic water had any effect on rat blood OCP.
5. Fetal analysis revealed that offspring of mother rats that had been watered with electrolysed water were smaller than those, whose mothers drank tap water.
6. Long bones of hind and front limbs of the fetuses from experimental groups (in particular – acidic) were noticeably shorter in comparison to the ones in control group.
7. In this model, ionised water had a negative effect on the development of embryos of mother rats which were given ionised (electrolysed) water.

Conflict of interest

The authors declare no conflict of interest.

References

- Abbasi, P. A., & Lazarovits, G. (2006). Effect of acidic electrolysed water on the viability of bacterial and fungal plant pathogens and on bacterial spot disease of tomato. *Canadian Journal of Microbiology*, 52, 915-923. <https://doi.org/10.1139/w06-048>
- Abdelhalim, M. I., Salaheldeen, Y. M., Idris, O. F., Abdelsalam, E. B., & Sabahelkher, M. K. (2016). Histopathological effect on different rat tissues induced by the trihalomethane-bromoform administered in drinking water. *Nova Journal of Medical and Biological Sciences*, 5, 1-7. <https://doi.org/10.20286/nova-jmbs-050280>
- Audickaitė, A., Stankevič, J., Šimčikas, V., Cesiulis, H., Jarašienė, R., Bukelskienė, V., Žalgevičienė, V., & Tutkuviene, J. (2014). Jonizuoto (elektrolizuoto) vandens įtaka žiurkių patelių fizinei būklei (eksperimentinis tyrimas) [The influence of ionized (electrolysed) water on physical status of female rats (experimental study)]. *Sveikatos mokslai [Health Sciences]*, 24, 46-51. <https://doi.org/10.5200/sm-hs.2014.024>
- Benmarhnia, T., Delpla, I., Schwarz, L., Rodriguez, M., & Levallois, P. (2018). Heterogeneity in the relationship between disinfection by-products in drinking water and cancer: a systematic review. *International Journal of Environmental Research and Public Health*, 15(5), 979. <https://doi.org/10.3390/ijerph15050979>
- Bui, V. N., Nguyen, K. V., Pham, N. T., Bui, A. N., Dao, T. D., Nguyen, H. T., Trinh, D. Q., Inui, K., Uchiumi, H., Ogawa, H., & Imai, K. (2017). Potential of electrolysed water for disinfection of foot-and-mouth disease virus. *Journal of Veterinary Medical Science*, 79, 726-729. <https://doi.org/10.1292/jvms.16-0614>
- Chen, M. J., Lin, C. H., Duh, J. M., Chou, W. S., & Hsu, H. T. (2011). Development of a multi-pathway probabilistic health risk assessment model for swimmers exposed to chloroform in indoor swimming pools. *Journal of Hazardous Materials*, 185, 1037-1044. <https://doi.org/10.1016/j.jhazmat.2010.10.011>
- Dawson-Hughes, B. (2016). Acid-base balance of the diet: implications for bone. In C. M. Weaver, R. M. Daly, & H. A. Bischoff-Ferrari (Eds.), *Nutritional influences on bone health (9th International Symposium)*. Cham: Springer International Publishing Switzerland. https://doi.org/10.1007/978-3-319-32417-3_9
- El-Fiky, N. K. (2002). The influence of water pH on the embryonic development of grass carp, *Ctenopharyngodon idell*. *Egyptian Journal of Aquatic Biology and Fisheries*, 6, 233-261. <https://doi.org/10.21608/ejabf.2002.1760>
- Fadel, R. A., & Persaud, T. V. (1993). Ossification of the vertebral column in the offspring of rats exposed to alcohol, acetaldehyde and caffeine. *Experimental and Toxicologic Pathology*, 45, 51-54. [https://doi.org/10.1016/S0940-2993\(11\)80456-2](https://doi.org/10.1016/S0940-2993(11)80456-2)
- Fadel, R. A., Sequeira, R. P., Abu-Hijleh, M. F., Obeidat, M., & Salem, A. H. (2012). Effect of prenatal administration of therapeutic doses of topiramate on ossification of ribs and vertebrae in rat fetuses. *Romanian Journal of Morphology and Embryology*, 53, 321-327.
- Fenton, T. R., Eliasziw, M., Lyon, A. W., Tough, S. C., & Hanley, D. A. (2008). Meta-analysis of the quantity of calcium excretion associated with the net acid excretion of the modern diet under the acid-ash diet hypothesis. *The American Journal of Clinical Nutrition*, 88, 1159-1166. <https://doi.org/10.1093/ajcn/88.4.1159>
- Fenton, T. R., & Huang, T. (2016). Systematic review of the association between dietary acid load, alkaline water and cancer. *BMJ Open*, 6, 1-5. <https://doi.org/10.1136/bmjopen-2015-010438>
- Fenton, C. J., Fenton, T. R., & Huang, T. (2017). Further evidence of no association between dietary acid load and disease. *The Journal of Nutrition*, 147(2), 272. <https://doi.org/10.3945/jn.116.242107>
- Flem, B., Reimann, C., Birke, M., Banks, D., & Filzmoser, P. (2015). Inorganic chemical quality of European tap-water: 2. Geographical distribution. *Applied Geochemistry*, 59, 211-224. <https://doi.org/10.1016/j.apgeochem.2015.01.016>
- Florentin, A., Hautemaniere, A., & Hartemann, P. (2011). Health effects of disinfection by-products in chlorinated swimming pools. *International Journal of Hygiene and Environmental Health*, 214, 461-469. <https://doi.org/10.1016/j.ijheh.2011.07.012>
- Hayes, A. W. (1994). *Principles and methods of toxicology*. New York: Raven Press.
- Hrudey, S. E., Backer, L. C., Humpage, A. R., Krasner, S. W., Michaud, D. S., Moore, L. E., Singer, P. C., & Stanford, B. D. (2015). Evaluating evidence for association of human bladder cancer with drinking-water chlorination disinfection by-products. *Journal of Toxicology and Environmental Health, Part B*, 18, 213-241. <https://doi.org/10.1080/10937404.2015.1067661>
- Hu, H., Zhao, X., Ma, J., Shangguan, Y., Pan, Z., Chen, L., Zhang, X., & Wang, H. (2018). Prenatal nicotine exposure retards osteoclastogenesis and endochondral ossification in fetal long bones in rats. *Toxicology Letters*, 295, 249-255. <https://doi.org/10.1016/j.toxlet.2018.07.005>
- Ignacio, R. M. C., Joo, K.-B., & Lee, K.-J. (2012). Clinical effect and mechanism of alkaline reduced water. *Journal of Food and Drug Analysis*, 20, 394-397.
- Jensen, V. F. H., Molck, A.-M., Berthelsen, L. O., Andersen, L., Demozay, D., Renaut R., & Bogh, I. (2017). Gestational hypoglycaemia restricts foetal growth and skeletal ossification in the rat. *Journal of Pregnancy and Child Health*, 4, 1-11.
- Khubutiya, M. S., Evseev, A. K., Kolesnikov, V. A., Goldin, M. M., Davydov, A. D., Volkov, A. G., & Stepanov, A. (2010). Measurements of platinum electrode potential in blood and blood plasma and serum. *Russian Journal of Electrochemistry*, 46, 537-541. <https://doi.org/10.1134/S1023193510050071>
- Klimas, A., & Mališauskas, A. (2008). Boron, fluoride, strontium and lithium anomalies in fresh groundwater of Lithuania. *Geologija*, 50(2), 114-124. <https://doi.org/10.2478/v10056-008-0015-4>
- Koufman, J. A., & Johnston, N. (2012). Potential benefits of pH 8.8 alkaline drinking water as an adjunct in the treatment of reflux disease. *Annals of Otolaryngology, Rhinology & Laryngology*, 121, 431-434. <https://doi.org/10.1177/000348941212100702>
- Lack, T. (1999). Water and health in Europe: an overview. *BMJ*, 318(7199), 1678-1682. <https://doi.org/10.1136/bmj.318.7199.1678>
- Laucevičius, T. (2009). *Jonizuotas vanduo. Gyvenimas be ligų. [Ionised water. Life without diseases]*. Kaunas: Obuolys (Lithuania).
- Legay, C., Rodriguez, M. J., Sadiq, R., Sérodes, J. B., Levallois, P., & Proulx, F. (2011). Spatial variations of human health risk associated with exposure to chlorination by-products occurring in drinking water. *Journal of Environmental Management*, 92, 892-901. <https://doi.org/10.1016/j.jenvman.2010.10.056>
- LRP. (2016). *Lithuania's water resources and their protection receive attention in Davos*. Vilnius. Retrieved from <https://www.lrp.lt/en/press-centr/press-releases/lithuanias-water-resources-and-their-protection-receive-attention-in-davos/24575>.
- Magro, M., Corain, L., Ferro, S., Baratell, D., Bonaiuto, E., Terzo, M., Corraducci, V., Salmaso, L., & Vianello, F. (2016).

- Alkaline water and longevity: a murine study. *Evidence-Based Complementary and Alternative Medicine*, 2016, 1-6. <https://doi.org/10.1155/2016/3084126>
- Merne, M. E., Syrjanen, K. J., & Syrjanen, S. M. (2001). Systemic and local effects of long-term exposure to alkaline drinking water in rats. *International Journal of Experimental Pathology*, 82, 213-219. <https://doi.org/10.1111/j.1365-2613.2001.iep188.x>
- Mills, C. J., Bull, R. J., Cantor, K. P., Reif, J., Hrudey, S. E., & Huston, P. (1998). Workshop report. Health risks of drinking water chlorination by-products: report of an expert working group. *Chronic Diseases in Canada*, 19, 91-102.
- Mousa, H. A. (2016). Health effects of alkaline diet and water, reduction of digestive-tract bacterial load, and earthing. *Alternative Therapies in Health and Medicine*, 22, 24-33.
- Ornoy, A., Rand, S. B., & Bischitz, N. (2010). Hyperglycemia and hypoxia are interrelated in their teratogenic mechanism: studies on cultured rat embryos. *Birth Defects Research. Part B, Developmental and Reproductive Toxicology*, 89, 106-115. <https://doi.org/10.1002/bdrb.20230>
- Osman, K. T. (2018). *Management of soil problems*. Cham: Springer International Publishing AG. <https://doi.org/10.1007/978-3-319-75527-4>
- Rashid, M. M., Al Mesfer, M. K., Naseem, H., & Danish, M. (2015). Hydrogen production by water electrolysis: a review of alkaline water electrolysis, PEM water electrolysis and high temperature water electrolysis. *International Journal of Engineering and Advanced Technology*, 4, 80-93.
- Rauch, F., & Schoenau, E. (2001). The developing bone: slave or master of its cells and molecules? *Pediatric Research*, 50, 309-314. <https://doi.org/10.1203/00006450-200109000-00003>
- Santos, B. A. F., Serenário, M. E. D., Pinto, D. L. M. F., Simões, T. A., Malafaia, A. M. S., & Bueno, A. H. S. (2019). Evaluation of micro-crack incidence and their influence on the corrosion resistance of steel coated with different chromium thicknesses. *Revista Virtual de Química*, 11, 264-274. <https://doi.org/10.21577/1984-6835.20190019>
- Shirahata, S., Kabayama, S., Nakano, M., Miura, T., Kusumoto, K., Gotoh, M., Hayashi, H., Otsubo, K., Morisawa, S., & Katakura, Y. (1997). Electrolysed-reduced water scavenges active oxygen species and protects DNA from oxidative damage. *Biochemical and Biophysical Research Communications*, 234, 269-274. <https://doi.org/10.1006/bbrc.1997.6622>
- Shirahata, S., Hamasaki, T., & Teruya, K. (2012). Advanced research on the health benefit of reduced water. *Trends in Food Science & Technology*, 23, 124-131. <https://doi.org/10.1016/j.tifs.2011.10.009>
- Sun, J. L., Zhang, S. K., Chen, J. Y., & Han, B. Z. (2012). Efficacy of acidic and basic electrolysed water in eradicating *Staphylococcus aureus* biofilm. *Canadian Journal of Microbiology*, 58, 448-454. <https://doi.org/10.1139/w2012-005>
- Villanueva, C. M., Cordier, S., Font-Ribera, L., Salas, L. A., & Levallois, P. (2015). Overview of disinfection by-products and associated health effects. *Current Environmental Health Reports*, 2, 107-115. <https://doi.org/10.1007/s40572-014-0032-x>
- Vilnius vandens. (2016). *Water quality*. Retrieved from <https://www.vv.lt/en/quality/#>
- Volker, J., & Borchardt, D. (2019). Drinking water quality at risk: a European perspective. In M. Schroter, A. Bonn, S. Klotz, R. Seppelt, & Baessler, C. (Eds.), *Atlas of ecosystem services. Drivers, risks, and societal responses*. Cham: Springer International Publishing AG.
- Vormann, J., & Goedecke, T. (2002). Latent acidosis: overacidification as a cause of chronic diseases. *Journal Suisse de Médecine Globale*, 14, 90-96.
- Watanabe, T., Kishikawa, Y., & Shirai, W. (1997). Influence of alkaline ionised water on rat erythrocyte hexokinase activity and myocardium. *Journal of Toxicological Sciences*, 22, 141-152. https://doi.org/10.2131/jts.22.2_141
- Watanabe, T., & Kishikawa, Y. (1998a). Degradation of myocardial myosin and creatine kinase in rats given alkaline ionised water. *Journal of Veterinary Medical Science*, 60, 245-250. <https://doi.org/10.1292/jvms.60.245>
- Watanabe, T., Shirai, W., Pan, I., Fukuda, Y., Murasugi, E., Sato, T., Kamata, H., & Uwatoko, K. (1998b). Histopathological influence of alkaline ionised water on myocardial muscle of mother rats. *Journal of Toxicological Sciences*, 23, 411-417. https://doi.org/10.2131/jts.23.5_411
- Weidman, J., Holsworth, R. E., Brossman, B., Cho, G. J., Cyr, J. S., & Fridman, G. (2016). Effect of electrolysed high-pH alkaline water on blood viscosity in healthy adults. *Journal of the International Society of Sports Nutrition*, 13(45), 1-13. <https://doi.org/10.1186/s12970-016-0153-8>
- Weldeslassie, T., Naz, H., Singh, B., & Oves, M. (2018). Chemical contaminants for soil, air and aquatic ecosystem. In M. Oves, M. Z. Khan, & I. M. I. Ismail (Eds.), *Modern age environmental problems and their remediation*. Cham: Springer International Publishing AG. https://doi.org/10.1007/978-3-319-64501-8_1
- World Health Organization. (2005). *Nutrients in drinking water*. Geneva. Retrieved from http://www.who.int/water_sanitation_health/dwq/nutrientsindw.pdf?ua=1
- Xu, H., Akesson, A., Orsini, N., Hakansson, N., Wolk, A., & Carrero, J. J. (2016). Modest U-shaped association between dietary acid load and risk of all-cause and cardiovascular mortality in adults. *The Journal of Nutrition*, 146, 1580-1585. <https://doi.org/10.3945/jn.116.231019>
- Xue, J., Shang, G., Tanaka, Y., Saihara, Y., Hou, L., Velasquez, N., Liu, W., & Lu, Y. (2014). Dose-dependent inhibition of gastric injury by hydrogen in alkaline electrolyzed drinking water. *BMC Complementary and Alternative Medicine*, 14, (81), 1-9. <https://doi.org/10.1186/1472-6882-14-81>
- Zukiene, J., Zalgevičienė, V., & Rizgeliene, R. (2003). The influence of azathioprine on the osteogenesis of the limbs. *Medicina*, 39, 584-588.